

Neutrino Masses and SO(10) SUSY GUTs

Stuart Raby^{a,*}

^aDepartment of Physics, The Ohio State University, 174 W. 18 th Ave, Columbus, Ohio 43210, USA

In this talk I discuss the qualitative features of an SO(10) SUSY GUT with an $SU(2) \times U(1)^n$ family symmetry. I then describe the global fit of this theory to precision electroweak data [including charged fermion masses and mixing angles]. Finally, the predictions of the model for neutrino masses and mixing are discussed. The most predictive version of the model naturally fits atmospheric neutrino data with maximal $\nu_\mu \rightarrow \nu_\tau$ oscillations and solar neutrino data with small mixing angle MSW $\nu_e \rightarrow \nu_{sterile}$ oscillations.

1. The Steps in Model Building

Before I discuss a particular model, let me first describe the steps one must take when constructing a candidate theory of nature.

• List the problems

The initial step in model building requires one to analyze the experimental data and list the problems/questions left unresolved in the Standard Model. For example, we have the phenomenological/theoretical problems

1. Gauge hierarchy – $M_Z/M_{Planck} \sim 10^{-17}$
2. Pattern of fermion masses and mixing
3. Absence of large flavor violation

• List the problem solving mechanisms

The next step is to obtain a list of possible mechanisms for solving the above problems. Each mechanism is capable of solving at least one (but preferably more than one) problem. Note, many known mechanisms are mutually exclusive.

• Develop a theoretical framework

*On leave at Universitat Bonn, Physikalisches Institut, Nussallee 12, 53115 Bonn, Germany. The work presented in this talk was done in collaboration with T. Blažek and K. Tobe. Partial support for this work came from an Alexander von Humboldt Fellowship and a DOE grant DOE/ER/01545.

Finally, one develops a theoretical framework incorporating a maximal set of *mutually consistent* mechanisms.

Thus prior to presenting any specific model, let us consider the list of [**mechanisms/problems**] we hope to incorporate into our model.

- SUSY - *a mechanism* for solving the gauge hierarchy problem.
- GUTs - *for* explaining charge quantization; family structure and reducing the number of fundamental parameters.
- Textures - *for* obtaining the pattern of fermion masses and mixing.
- Family Symmetry - *for* explaining textures; the hierarchy of family masses and solving the flavor problem.

2. The Model

The model is an SO_{10} SUSY GUT with an $SU_2 \times U_1^n$ family symmetry[1,3,4]. The three families of quarks and leptons belong to the spinor representations 16_a , 16_3 where $a = 1, 2$ is an SU_2 family index.

Effective 3×3 Yukawa matrices are obtained at the GUT scale after spontaneously breaking the family symmetry and integrating out states with mass (M , M_χ) above M_G . Schematically they have the form

$$\begin{pmatrix} 0 & \epsilon' D & 0 \\ -\epsilon' D & \epsilon C & \epsilon B \\ 0 & \epsilon B & A \end{pmatrix}$$

where the small parameters

$$\epsilon = \frac{\langle \phi^2 \rangle \langle 45 \rangle}{MM_\chi} \approx \frac{\langle S^{22} \rangle \langle 45 \rangle}{MM_\chi}$$

$$\epsilon' = \frac{\langle A^{12} \rangle}{M}$$

are proportional to the family symmetry breaking vacuum values of the $SO(10)$ singlet fields ϕ^a , $S^{ab} = S^{ba}$, $A^{ab} = -A^{ba}$ and the SO_{10} adjoint field 45.

2.1. Features of the Model

1. Family symmetry breaking above the GUT scale generates the hierarchy of fermion masses. In the symmetric limit, only the third generation can have mass. The mass hierarchy for the second and first generations is determined by the family symmetry breaking ratios (ϵ , ϵ').

$$\begin{array}{ccccc} SU_2 \times U_1 & \longrightarrow & U_1 & \longrightarrow & \text{nothing} \\ & & \epsilon & & \epsilon' \end{array}$$

3rd family $\gg$ 2nd family $\gg$ 1st family

2. Some well known patterns for fermion masses are satisfied approximately in the model. For example, the Georgi - Jarlskog relation $m_s \sim \frac{1}{3}m_\mu$ $m_d \sim 3m_e$ @ M_G results from the vev of the adjoint $\langle 45 \rangle = (B-L)M_G$ in the 22 matrix element, where $B-L$ is baryon number minus lepton number. The same adjoint is responsible for Higgs doublet-triplet splitting.
3. Yukawa coupling unification for the third generation is well respected by the data. In particular, the relation $\lambda_t = \lambda_b = \lambda_\tau = \lambda_{\nu_\tau} = A$ @ M_G gives the result $m_t \sim 170 \pm 20$ GeV.
4. Gauge coupling unification is well respected by the data.

5. The SU_2 family symmetry naturally suppresses flavor violation such as $\mu \rightarrow e\gamma$.

6. Finally the model has **only 9 arbitrary Yukawa parameters to fit 13 charged fermion masses and mixing angles**. Hence there are 4 non-trivial predictions.

2.2. Global χ^2 fit to precision electroweak data[3,4]

We fit the data using a global χ^2 analysis discussed originally in [2]. This analysis includes 2 loop RG running from $M_G \rightarrow M_Z$ with the following boundary conditions at M_G .

- universal squark and slepton masses - m_0 ,
- universal gaugino masses - $M_{1/2}$,
- non-universal H_u , H_d masses,
- $\epsilon_3 (\equiv \frac{\alpha_3 - \alpha_G}{\alpha_G})$ parameterizes the effect of one loop threshold corrections to the unification of the three gauge coupling constants at M_G . Note the GUT scale is defined as the scale where $\alpha_1 = \alpha_2$.

We also include one loop threshold corrections at M_Z to the Z and W masses and the leading corrections to the down-type quark and charged lepton masses proportional to $\tan\beta$. We use 3 loop QCD + 1 loop QED RG running below M_Z . We demand a consistent electroweak symmetry breaking solution using the one loop improved Higgs potential, including m_t^4 corrections. Finally we construct a χ^2 **function including 20 low energy observables**. The result of this fit is given in the following table (for details, see[3,4]). Note, the observables in bold script have the largest pulls.

The parameters m_0 , $M_{1/2}$ and μ were fixed. All others were varied using Minuit. The following are the parameters determined at M_G for the χ^2 fit in the table.

$$\begin{aligned} (1/\alpha_G, M_G, \epsilon_3) &= (24.52, 3.03 \cdot 10^{16} \text{ GeV}, -4.06\%), \\ (\lambda, r, \sigma, \epsilon, \rho, \epsilon') &= (0.79, 12.4, 0.84, 0.011, 0.043, 0.0031), \\ (\Phi_\sigma, \Phi_\epsilon, \Phi_\rho) &= (0.73, -1.21, 3.72)\text{rad}, \\ (m_0, M_{1/2}, A_0, \mu(M_Z)) &= (1000, 300, -1431, \end{aligned}$$

110) GeV,
 $((m_{H_d}/m_0)^2, (m_{H_u}/m_0)^2, \tan\beta) = (2.23, 1.66, 53.7)$

Table 1
 Fit to charged fermion masses and mixing angles.

Observable	Data(σ) (masses)	Theory in GeV)
M_Z	91.187 (0.091)	91.17
M_W	80.388 (0.080)	80.39
$G_\mu \cdot 10^5$	1.1664 (0.0012)	1.166
α_{EM}^{-1}	137.04 (0.14)	137.0
$\alpha_s(\mathbf{M}_Z)$	0.1190 (0.003)	0.1174
$\rho_{new} \cdot 10^3$	-0.20 (1.1)	+0.314
M_t	173.8 (5.0)	174.9
$m_b(M_b)$	4.260 (0.11)	4.331
$M_b - M_c$	3.400 (0.2)	3.426
$\mathbf{m}_s$	0.180 (0.050)	0.147
m_d/m_s	0.050 (0.015)	0.0589
Q^{-2}	0.00203 (0.00020)	0.00201
M_τ	1.777 (0.0018)	1.777
M_μ	0.10566 (0.00011)	.1057
$M_e \cdot 10^3$	0.5110 (0.00051)	0.5110
V_{us}	0.2205 (0.0026)	0.2205
V_{cb}	0.03920 (0.0030)	0.0403
$\mathbf{V}_{ub}/\mathbf{V}_{cb}$	0.0800 (0.02)	0.0691
$\hat{B}_K$	0.860 (0.08)	0.870
$B(b \rightarrow s\gamma) \cdot 10^4$	3.000 (0.47)	2.992
TOTAL χ^2	2.26	

Of course once the data is fit, there are other predictions of the theory. As an example we obtain the CP violating parameters measured in B decays $\sin 2\alpha = 0.74$, $\sin 2\beta = 0.54$, $\sin \gamma = 0.99$ and the SUSY contribution to the anomalous magnetic moment of the muon $a_\mu^{SUSY} = 40 \times 10^{-10}$. We also obtain the complete spectrum of squark, slepton, Higgs, and gaugino masses.

2.3. Neutrinos in Standard Model

Let's now consider neutrino masses and mixing. In the Standard Model neutrinos masses are obtained by introducing three sterile neutrinos $\bar{\nu}$. The neutrino masses are then given by the expression $l U_e^\dagger \lambda_\nu \bar{\nu} H_u + \frac{1}{2} \bar{\nu}^T M \bar{\nu}$ where the sec-

ond term is an invariant mass term for the sterile neutrinos. Note, the charged leptons in the weak doublet l are mass eigenstates and U_e is the unitary matrix used to diagonalize the charged lepton mass matrix. The mass scale M is not constrained and thus could be taken much larger than M_Z . As a consequence, the heavy sterile neutrinos may be integrated out of the theory, leaving behind the effective 3 x 3 neutrino mass matrix given by $m_\nu^{eff} = U_e^\dagger m_\nu M^{-1} m_\nu^T U_e^*$.

2.4. Neutrinos in SO(10)

A Dirac neutrino mass matrix m_ν given by $\nu m_\nu \bar{\nu}$ (with $\nu, \bar{\nu}$ in the 16) has already been fixed by the fits for charged fermions.

The most economical way of obtaining the see-saw (without introducing higher dimension operators) is by adding three SO(10) singlets N into the superpotential as follows - $16 \mathbf{\bar{16}} N + \frac{1}{2} N \mathbf{M}_N N$ where the field $\mathbf{\bar{16}}$ obtains a vev in the right-handed neutrino direction and $\mathbf{M}_N$ is a mass term. Both are assumed to be of order M_G . This gives the effective mass terms $N \mathbf{V} \bar{\nu} + \frac{1}{2} N \mathbf{M}_N N$. We then obtain the 9 x 9 neutrino mass matrix at M_G given by

$$\begin{pmatrix} \nu & \bar{\nu} & N \end{pmatrix} \begin{pmatrix} 0 & m_\nu & 0 \\ m_\nu^T & 0 & V \\ 0 & V^T & M_N \end{pmatrix}$$

Upon integrating out the massive states $\bar{\nu}, N$ we obtain the effective 3 x 3 neutrino mass matrix - $m_\nu^{eff} = U_e^\dagger m_\nu (V^T)^{-1} M_N V^{-1} m_\nu^T U_e^*$. Now consider the particular SO(10) theory defined earlier.

2.5. $SO_{10} \times SU_2 \times U_1^n$ and Neutrino Masses[3]

The three SO(10) singlets are given by N_a, N_3 with $a = 1, 2$. The superpotential below is the most general one consistent with the family symmetries.

$$W \supset \mathbf{\bar{16}} (N_a \chi^a + N_3 16_3) + \frac{1}{2} N_a N_b S^a b + N_a N_3 \phi^a$$

We now see that the effective mass matrix $\mathbf{M}_N$ discussed above is constrained by the family sym-

metries and has the form

$$M_N = \begin{pmatrix} 0 & 0 & 0 \\ 0 & S^{22} & \phi^2 \\ 0 & \phi^2 & 0 \end{pmatrix}$$

We have analyzed several different possibilities[3]. In the simplest case, we have only three light active neutrinos. The effective 3 x 3 neutrino mass matrix is given by

$$m_\nu^{eff} = m' U_e^\dagger \begin{pmatrix} 0 & 0 & 0 \\ 0 & b & 1 \\ 0 & 1 & 0 \end{pmatrix} U_e^*$$

There are only two free parameters in this matrix given by the overall scale m' and b . Both may be taken real without affecting the observable neutrino masses and mixing angles. The zero in the 33 term is due to an accidental cancellation. However as a result we naturally obtain maximal $\nu_\mu \rightarrow \nu_\tau$ mixing for any value of $b \sim \frac{\langle S^{22} \rangle}{\langle \phi^2 \rangle} \leq 1$. Given this neutrino mass matrix, we find that we can fit both *atmospheric* and *LSND* but **NOT** *atmospheric* and *solar* neutrino oscillations!!

If we want to fit both atmospheric and solar neutrino oscillations we must add a *light sterile neutrino*. I'll describe how we do this next, but in the meantime let me just describe the results. We now find a solution to *atmospheric* neutrino oscillations with maximal $\nu_\mu \rightarrow \nu_\tau$ mixing and a fit to *solar* data given by the **SMA MSW** solution with $\nu_e \rightarrow \nu_{sterile}$ oscillations. Note, that even though we have four neutrinos, we are not able to fit LSND.

2.6. Sterile neutrinos in SUSY GUTs

We now argue that it is natural to have light sterile neutrinos in the MSSM. In principle any SUSY GUT could have many SO(10) singlets. The question is only whether they couple to the observable sector. Consider adding three such singlets $\bar{N}^a$, $\bar{N}^3$ with the dimensionful couplings given by

$$W \supset \mu' N_a \bar{N}^a + \mu_3 N_3 \bar{N}^3$$

The mass scales μ' , μ_3 are assumed to be of order the weak scale. Via standard steps we obtain the neutrino mass matrix at M_G

$$\begin{pmatrix} \nu & \bar{N} & \bar{\nu} & N \end{pmatrix} \begin{pmatrix} 0 & 0 & m_\nu & 0 \\ 0 & 0 & 0 & \tilde{\mu}^T \\ m_\nu^T & 0 & 0 & V \\ 0 & \tilde{\mu} & V^T & M_N \end{pmatrix}$$

with

$$\tilde{\mu} \equiv \begin{pmatrix} \mu' & 0 & 0 \\ 0 & \mu' & 0 \\ 0 & 0 & \mu_3 \end{pmatrix}.$$

The effective light neutrino mass matrix m_ν^{eff} is given by

$$m_\nu^{eff} = \tilde{U}_e^T \mathbf{m} \tilde{U}_e$$

where

$\mathbf{m} =$

$$\begin{pmatrix} m_\nu (V^T)^{-1} M_N V^{-1} m_\nu^T & -m_\nu (V^T)^{-1} \tilde{\mu} \\ -\tilde{\mu}^T V^{-1} m_\nu^T & 0 \end{pmatrix}$$

and

$$\tilde{U}_e = \begin{pmatrix} U_e & 0 \\ 0 & 1 \end{pmatrix}.$$

Note we now have light sterile neutrinos and all we need do is assume that the scales μ' , μ_3 are of order M_Z . RECALL in the MSSM we have another example of a mass term in the superpotential, μ , defined via $W = \mu H_u H_d$ which must also be of order M_Z . What ever symmetries are needed to solve the μ problem in the MSSM could in principle also explain why our new "mu" parameters are also of order M_Z .

2.7. How robust is this neutrino solution?

The four neutrino solution found above cannot fit LSND data. It is important to investigate whether this is a true prediction of the theory. In order to study this we now assume the possibility of *non-minimal* family symmetry breaking vevs $\langle S^{11} \rangle = \kappa_1 \langle S^{22} \rangle$, $\langle S^{12} \rangle = \kappa_2 \langle S^{22} \rangle$ with the parameters $\kappa_1, \kappa_2 \ll 1$ [4]. In the model presented here, we are not able to find a dynamical argument why $\kappa_1 = \kappa_2 \equiv 0$ should be an exact relation. Moreover this extension of the original model has little effect on charged fermion fits for

sufficiently small values of κ_1, κ_2 . However we now find several *new solutions* given below. In each solution the atmospheric oscillation data is described by maximal $\nu_\mu \rightarrow \nu_\tau$ mixing.

We have a 3 ν solution with three possibilities for solar neutrinos – either SMA MSW; LMA MSW, or Vacuum $\nu_e \rightarrow \nu_{active}$ oscillations. In addition we have 4 ν and 5 ν solutions with solar data given by SMA MSW $\nu_e \rightarrow \nu_{sterile}$ oscillations and LSND given by $\nu_e \rightarrow \nu_\mu$.

3. Conclusions

The theoretical framework presented here provides a compact and precise description of low energy data. With regards to neutrino masses and mixing we have two possibilities –

- In a “predictive” theory (with sufficient symmetry to enforce a small number of arbitrary parameters) neutrino masses are very constrained. In this case, charged fermion masses and mixing angles can strongly constrain the neutrino sector.

For example, an $SO_{10} \times D_3 \times Z_M$ model provides a natural framework for the case of minimal family symmetry breaking vevs [5].

- If there are however many free parameters, then neutrino masses and mixing are poorly constrained by the charged fermion sector.

Finally we have not presented a complete theory here. There are still many unresolved problems : SUSY breaking, GUT breaking, family symmetry breaking, μ problem, strong CP problem.

REFERENCES

1. R. Barbieri, L.J. Hall, S. Raby and A. Romanino, Nucl. Phys. B493 (1997) 3.
2. T. Blažek, M. Carena, S. Raby and C.E.M. Wagner, Phys. Rev. D56 (1997) 6919.
3. T. Blažek, S. Raby and K. Tobe, Phys. Rev. D60 (1999) 113001.
4. T. Blažek, S. Raby and K. Tobe, Phys. Rev. D62 (2000) 055001.
5. R. Dermíšek and S. Raby, Phys. Rev. D62 (2000) 015007.